



**HAL**  
open science

## Chemical multi-contamination drives benthic prokaryotic diversity in the anthropized Toulon Bay

Benjamin Misson, Cédric Garnier, Béatrice Lauga, Duc Huy Dang, Jean-François Ghiglione, Jean-Ulrich Mullot, Robert Duran, Olivier Pringault

► **To cite this version:**

Benjamin Misson, Cédric Garnier, Béatrice Lauga, Duc Huy Dang, Jean-François Ghiglione, et al.. Chemical multi-contamination drives benthic prokaryotic diversity in the anthropized Toulon Bay. Science of the Total Environment, 2016, 556, pp.319 - 329. 10.1016/j.scitotenv.2016.02.038 . hal-01826037

**HAL Id: hal-01826037**

**<https://univ-tln.hal.science/hal-01826037v1>**

Submitted on 26 Jan 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Chemical multi-contamination drives benthic prokaryotic diversity in the anthropized Toulon Bay

Benjamin Misson<sup>a</sup>, Cédric Garnier<sup>a</sup>, Béatrice Lauga<sup>b</sup>, Duc Huy Dang<sup>a,1</sup>, Jean-François Ghiglione<sup>d</sup>, Jean-Ulrich Mullot<sup>e</sup>, Robert Duran<sup>b</sup> and Olivier Pringault<sup>f,g</sup>

<sup>a</sup> PROTEE, EA 3819, Université de Toulon, CS 60584, 83041 Toulon Cedex 9, France

<sup>b</sup> Equipe Environnement et Microbiologie, Melody group, Université de Pau et des Pays de l'Adour, IPREM, UMR CNRS 5254, BP 11055, F-64013 Pau cedex, France

<sup>d</sup> Sorbonne Universités, UPMC Univ Paris 06, CNRS, Laboratoire d'Océanographie Microbienne (LOMIC) UMR 7621, Observatoire Océanologique, F-66650, Banyuls/mer, France;

<sup>e</sup> LASEM de Toulon, Base Navale Toulon, BP 61, 83800 Toulon Cedex 9, France

<sup>f</sup> MARBEC, UMR 9190, CNRS IRD IFREMER Université Montpellier 2, F-34095 Montpellier, France

<sup>g</sup> Laboratoire de Biosurveillance de l'Environnement, Faculté des Sciences de Bizerte, Zarzouna 7021, Tunisie

<sup>1</sup> Present address: Water Quality Center, Trent University, 1600 West Bank Drive, Peterborough, ON, Canada K9J 7B8

Corresponding author: [mission@univ-tln.fr](mailto:mission@univ-tln.fr)

## ABSTRACT

1  
2  
3  
4 Investigating the impact of human activities on marine coastal ecosystems remains difficult  
5  
6 because of the co-occurrence of numerous natural and human-induced gradients. Our aims  
7  
8 were (i) to evaluate the links between the chemical environment as a whole and microbial  
9  
10 diversity in the benthic compartment, and (ii) to compare the contributions of anthropogenic  
11  
12 and natural chemical gradients to microbial diversity shifts. We studied surface sediments  
13  
14 from 54 sampling sites in the semi-enclosed Toulon Bay (NW Mediterranean) exposed to  
15  
16 high anthropogenic pressure. Previously published chemical data were completed by new  
17  
18 measurements, resulting in an in depth geochemical characterization by 29 representative  
19  
20 environmental variables. Bacterial and archaeal diversity was assessed by terminal  
21  
22 restriction fragment length polymorphism profiling on a selection of samples distributed along  
23  
24 chemical gradients. Multivariate statistical analyses explained from 45% to 80% of the spatial  
25  
26 variation in microbial diversity, considering only the chemical variables. A selection of trace  
27  
28 metals of anthropogenic origin appeared to be strong structural factors for both bacterial and  
29  
30 archaeal communities. Bacterial terminal restriction fragment (T-RF) richness correlated  
31  
32 strongly with both anthropogenic and natural chemical gradients, whereas archaeal T-RF  
33  
34 richness demonstrated fewer links with chemical variables. No significant decrease in  
35  
36 diversity was evidenced in relation to chemical contamination, suggesting a high adaptive  
37  
38 potential of benthic microbial communities in Toulon Bay.  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

51 **Key words:** microbial ecotoxicology, multi-contamination, prokaryotic diversity, marine  
52 sediment  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 1. INTRODUCTION

1  
2 Marine coastal areas represent ecotones between continental and marine environments.  
3  
4 Strong and natural coastal gradients (*e.g.* salinity, water inflows, bottom topography,  
5  
6 sedimentation pattern, turbidity and nutrients) are known to shape the diversity of several  
7  
8 organisms, ranging from fishes (Winemiller and Leslie, 1992) to bacteria (Ortega-Retuerta et  
9  
10 al., 2013; Troussellier et al., 2002), including invertebrates (Waterkeyn et al., 2008). As a  
11  
12 result, the functions of coastal ecosystems are largely driven by natural gradients (Sklar and  
13  
14 Browder, 1998). Since coastal areas are of great interest to humans, they also face  
15  
16 increasing human-induced chemical contamination resulting from watershed drainage,  
17  
18 domestic and industrial waste waters and other direct inputs, such as biocides released by  
19  
20 anti-fouling paints or fuel residues from shipping (*e.g.* Colombo et al., 2006; Daskalakis and  
21  
22 Oconnor, 1995; Levin et al., 2009; Oursel et al., 2014, 2013). As a result, marine coastal  
23  
24 areas are characterized by a complex set of chemical factors potentially affecting living  
25  
26 organisms. Consequently, marine coastal ecosystems represent one of the most challenging  
27  
28 environments for understanding the links between chemical composition and biological  
29  
30 diversity.  
31  
32  
33  
34

35 Benthic archaea and bacteria are key microbial players in ecosystem functioning, because of  
36  
37 their major roles in diverse biogeochemical cycles, particularly through carbon and nitrogen  
38  
39 mineralization (Francis et al., 2005; Jørgensen, 2000) and toxic metals transformation, either  
40  
41 directly (*e.g.* mercury methylation, Parks et al., 2013) or indirectly (*e.g.* formation of sulfide  
42  
43 compounds after sulfide release by sulfate reduction, Jørgensen, 2000). However, little is  
44  
45 known about the links between prokaryotes diversity and the global chemical composition of  
46  
47 the sediment, including both the natural composition and anthropogenic chemical  
48  
49 contaminants, in marine coastal areas. It was evidenced that early diagenesis, which is  
50  
51 controlled by microbial activity, induced modifications to the characteristics of sedimentary  
52  
53 organic matter (*e.g.* Burdige, 2007; Dang et al., 2014a) but also influenced the mobility of  
54  
55 trace elements, such as trace metals/metalloids (*e.g.* Dang et al., 2015a, 2014b; Gendron et  
56  
57 al., 1986), with significant seasonal variations. Moreover, the influence of microbial activity on  
58  
59  
60  
61  
62  
63  
64  
65

1 metal speciation has been demonstrated, particularly for mercury speciation (Cossa et al.,  
2 2014; Duran et al., 2008) and organic tin degradation (Filipkowska et al., 2014; Pougnet et  
3 al., 2014) in coastal sediments. Our understanding of the influence of benthic microbial  
4 activity on the chemical composition of anthropized coastal sediment is thus progressing  
5 well. Nevertheless, the impact of the global chemical composition of sediment on benthic  
6 microbial communities is poorly documented. Some bacteria and microalgae were shown to  
7 resist or even benefit from high inorganic or organic contaminant concentrations (e.g.  
8 Edwards et al., 2013; Gauthier et al., 1992; Louati et al., 2013; Rensing et al., 2000).  
9 However, chemical pollution can affect the diversity and functions of prokaryotic communities  
10 (Abell et al., 2014; Scott et al., 2014), although the great adaptive skills of prokaryotes after  
11 long term exposure to chemical contaminants (Gillan et al., 2014) can lead us to observe no  
12 significant change in diversity (Besaury et al., 2014; Gillan et al., 2005), not even higher  
13 diversities (Bengtsson et al., 2013) in response to chemical pollution. Such influences result  
14 in spatial variability of microbial abundance and/or community composition in the  
15 environment, which can alter ecosystem functioning along the contamination gradients (Abell  
16 et al., 2014; Scott et al., 2014). Thus, in marine coastal ecosystems, the relative contributions  
17 of natural abiotic gradients and anthropogenic and multiple contamination gradients to the  
18 spatial variability of microbial diversity need to be further explored in order to better  
19 apprehend the ecological consequences of human-induced environmental changes.

20 The present study was held at Toulon Bay, a semi-enclosed coastal ecosystem experiencing  
21 very strong and multiple contaminations by metals, metalloids, organometallic and organic  
22 pollutants (Cossa et al., 2014; Dang et al., 2015a, 2014b; Pougnet et al., 2014; Tessier et al.,  
23 2011). While this ecosystem experiences multiple contamination gradients of several orders  
24 of magnitude, its relatively small size greatly limits other abiotic variations such as climate  
25 related parameters, hydrology and mean sediment composition (excluding pollutants). An in-  
26 depth geochemical characterization of surface sediments and the genetic fingerprinting of  
27 archaeal and bacterial benthic communities sampled at numerous stations in Toulon Bay  
28 allowed us (i) to address the impact of the global chemical environment and then (ii) to

compare the respective influence of a wide range of both natural and anthropogenic gradients on both local microbial genetic richness and spatial variations in microbial community structure.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 2. MATERIALS AND METHODS

### 2.1. Study site and sediment sampling

Sediments samples were collected at different locations of Toulon Bay, on the northwest Mediterranean coast. Its semi-close morphology associated with micro-tidal conditions and intermittent freshwater inputs (Nicolau et al., 2012), leads to relatively long water residence times (Dufresne et al., 2014). Consequently, numerous historic (e.g. World War II) and present (1<sup>st</sup> French navy harbour, nautical traffic, tourism, urbanism, aquaculture, etc.) anthropogenic activities have resulted in a strong multi-contamination of the sedimentary compartment (Cossa et al., 2014; Dang et al., 2015a, 2014b; Pougnet et al., 2014; Tessier et al., 2011) and also of the water column (Jean et al., 2012, Dang et al., 2015b). A map of Toulon Bay including the location of the main towns and anthropogenic activities, is presented in figure 1A. The bay is separated into two basins by a sea wall: Little Bay (LiB) on the northwestern side and Large Bay (LaB) on the southeastern part. Sediments were collected using an interface corer (equipped with 10-cm diameter and 1-m long Plexiglass tube, handled from a navy boat or divers) as previously described (Tessier et al., 2011) to collect sediments at 54 stations covering the whole bay (figure 1).

### 2.2. Chemical characterization of the sediments

We compiled previously published data on the chemical characterization of the surface sediments, except for the organic pollutants (polycyclic aromatic hydrocarbons (PAHs), polychloro biphenylated (PCBs)). Briefly, the obtained surface sediments (0-5 cm) were analysed for (1) major constituents (major elements, particulate organic and inorganic carbon (POC, PIC), total phosphorus) and minor/trace elements (Dang et al., 2015a, 2014a, 2014b; Tessier et al., 2011); (2) methyl mercury (MMHg) (Cossa et al., 2014); and (3) tin and butyl-tin species (Pougnet et al., 2014). Full description of samples treatment and chemical analysis can be found in the cited references. Briefly, sediments were deep-frozen, freeze-dried, 2-mm sieved and kept deep frozen (-18°C) until analysis. POC and PIC contents were determined using a TOC-V<sub>CSH</sub> analyzer (Shimadzu) coupled with a SSM-5000A module

(Dang et al., 2014a; Tessier et al., 2011). Major elements and total phosphorus were measured by XRF (Philips PW 2400) equipped with a Rh-tube, on 50- $\mu$ m crushed (on carbide tungsten rings mill, NAEF model M02) sediments (Tessier et al., 2011). Minor/trace elements were determined by High Resolution Inductively Coupled Plasma Mass Spectrometer (HR ICP-MS, Element 2, Thermo Finnigan) after aqua regia and microwave (Anton Paar Multiwave 3000) digestion of sediments (Dang et al., 2014b, 2015a; Tessier et al., 2011). Hg contents were determined by an advanced mercury analyzer AMA 254 (Leco Corporation) (Tessier et al., 2011). MeHg was analysed after propylation and isotopic dilution and detection by gas chromatography-ICP-MS (GC/ICP-MS, GC-Focus with X-series, Thermo Electron) (Cossa et al., 2014). Butyl-tin species contents were determined by GC/ICP/MS (Thermo X7 Series II) after extraction of butyl-Sn species by acetic acid (Pougnnet et al., 2014). Moreover, 25 surface sediment samples from LiB (figure 1) were analysed for 17 PAHs (Acenaphthene, Anthracene, Benz[a]anthracene, Benzo[a]pyrene, Benzo[b]fluoranthene, Benzo[ghi]perylene, Benzo[k]fluoranthene, Chrysene, Dibenz[a,h]anthracene, Fluoranthene, Fluorene, Indeno[1,2,3-cd]pyrene, Naphthalene, Phenanthrene, Pyrene, 2-methylfluoranthene, 2-methylnaphthalene) and 8 PCBs (CB-28, CB-31, CB-52, CB-101, CB-118, CB-138, CB-153, CB-180), accordingly to the French standard AFNOR XP X33-012. Briefly, PAHs were extracted from 4 g of sediment using a mixture of hexane/acetone (50:50) assisted by ASE (ASE 200, Dionex), the obtained extract was then purified by solid-phase extraction (SPE, ASPEC XL, Gilson), concentrated (50°C under N<sub>2</sub> flux), adjusted to 5 mL with acetonitrile/water (50/50) and PAHs concentrations were analysed by HPLC (Shimadzu, Prominence UFLC) with UV and fluorescence detection. Concerning PCBs, they were extracted from 7 g of sediments by using a mixture of hexane/acetone (50:50) assisted by ASE (ASE 200, Dionex) in the presence of 1g of Cu powder. The obtained extract was then purified on SPE (ASPEC XL, Gilson), concentrated (50°C under N<sub>2</sub> flux), and adjusted to 1 mL with hexane. CBs concentrations were analysed by GC/MS (Thermoelectron, Polaris Q) in negative chemical ionization mode. The accuracy



1  
2 of these protocols was validated using certified material (MRC 1941b and MRC 1944 for  
3 PAHs and CBs, respectively).  
4  
5

### 6 **2.3. DNA extraction and prokaryotic fingerprinting**

7  
8 DNA was extracted from aliquots of lyophilized sediments (kept frozen at -20°C) from 38  
9 samples from among the 54 described above (figure 1), covering the whole range of main  
10 sediment characteristics and chemical contaminants levels. Extractions were performed  
11 using the Power Soil DNA Extraction Kit (Mobio Laboratories, Carlsbad, CA) following the  
12 manufacturer's recommendations. Terminal Restriction Fragment Length Polymorphism (T-  
13 RFLP) profiling was conducted on each sample after 16S rRNA gene amplification using  
14 primers 8F (Lane, 1991)/519R (Frank et al., 2013) labeled with FAM (5-carboxyfluorescein)  
15 and HEX (6-carboxy-1,4-dichloro-2',4', 5', 7'-tetrachlorofluorescein) fluorochromes  
16 respectively for Bacteria and 344F (Raskin et al., 1994)/915R (Stahl and Amann, 1991)  
17 labelled with FAM and HEX fluorochromes respectively for Archaea using AmpliTaq Gold®  
18 DNA Polymerase (Life technologies). For PCR amplification an initial denaturation step at  
19 94 °C for 5 min was followed by 35 cycles of 94 °C denaturation for 45 s, 56 °C annealing for  
20 45 s, and 72 °C extension for 1 min, and concluded with an extension at 72 °C for 10 min.  
21 PCR amplicons were cleaned using the GFX PCR DNA Purification kit (GE Fisher) and  
22 digested with the restriction endonucleases HinfI for Bacteria and TaqI for Archaea (37°C, 3  
23 h). T-RFLP profiles were obtained with a 3130xl Genetic Analyzer (Applied Biosystems)  
24 following the manufacturer's recommendations and analysed with GeneMapper®Software  
25 (Applied Biosystems). Using T-Rex software (Culman et al., 2009) raw data were normalized  
26 to account for differences in the amount of DNA loaded and terminal restriction fragments (T-  
27 RFs) assigned to produce the final matrix.  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

### 53 **2.4. Data analysis and statistics**

54  
55 The high spatial resolution maps allowed the establishment of surface-weighted average  
56 (SWA) trace metal/metalloids contents for Toulon Bay, *i.e.* the most representative  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 concentration values on the entire bay scale (Dang et al., 2015a; Larrose et al., 2010;  
2 Tessier et al., 2011). The kriging gridding method of Surfer software (9.9 version) with default  
3 parameters was used to generate contour map and Matlab software (7.8 version) was used  
4 for data processing.  
5  
6

7  
8 Using a principal component analysis (PCA) on 2 chemical characteristics (PIC, POC) and  
9 31 major/minor/trace elements (Ag, Al, As, Ba, Be, Bi, Ca, Cd, Co, Cr, Cs, Cu, Fe, Hg, K, Li,  
10 Mn, Mo, Ni, P, Pb, Rb, S, Sb, Sn, Sr, Ti, Tl, U, V and Zn) from 130 surface and deep  
11 sediments from Toulon Bay, Dang et al. (2015a) have evidenced 6 groups of elements: (1)  
12 the "marine" originated elements (Ca, PIC and Sr), (2) the "terrigenous" elements, mainly  
13 representing clays (Al, Be, Cs, Fe, K, Li and Rb), (3) "inorganic contaminants", i.e. elements  
14 mainly of anthropogenic origins (Ag, Ba, Bi, Cd, Cu, Hg, P, Pb, Sb, Sn, Ti and Zn), (4)  
15 inorganic trace elements having mainly "natural" origin, from terrigenous inputs (Co, Mo, Ni,  
16 Tl, U and V), (5) elements showing "mixed" anthropogenic and terrigenous origins (As, Cr,  
17 Mn and S), and (6) the POC. To illustrate these co-variations, the observed relationship  
18 between PIC and Ca, Rb vs. Al, Bi vs. Pb, Co vs. Ni and As vs. Cr in the surface sediments  
19 dataset considered in the present study are shown in Fig. SI-1. In order to reduce the  
20 number of chemical variables for further statistical analysis, the elements of "marine" origin  
21 were then only represented by Ca, the "terrigenous" elements by Al and Fe, and the trace  
22 elements of "natural" origin by Co and Ni, when POC, "inorganic contaminants" and trace  
23 elements of "mixed" origin were all taken into consideration. Then, a 7<sup>th</sup> group of  
24 "organometallic contaminants" was defined using: monomethylmercury (MMHg), tributyltin  
25 (TBT) and the sum of butyltin-species ( $\Sigma$ BT). Significant relationships observed between TBT  
26 and  $\Sigma$ BT, and MMHg and Hg (Fig. SI-2) were evidenced by Pougnet et al. (2014) and Cossa  
27 et al. (2014) and suggested the strong influence of microbial communities on the  
28 sedimentary dynamics of these species. Finally, an 8<sup>th</sup> group included the "organic  
29 contaminants" (PAHs and PCBs). Considering the significant correlations ( $R^2$  ranging from  
30 0.86 to 0.98 and from 0.95 to 0.99 for PAHs and PCBs respectively, data not shown)  
31 observed between each of the 17 measured PAHs and their sum ( $\Sigma$ PAH), and between each  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 of the 8 measured PCBs and their sum ( $\Sigma$ PCB), the present study only considered the  
2 following chemical variables:  $\Sigma$ PAH,  $\Sigma$ PCB, Fluoranthene and CB-153 (being the most  
3 abundant PAH and PCB, accounting on average for 16% and 29% (N=26) of the  $\Sigma$ PAH and  
4  $\Sigma$ PCB, respectively; figure SI-3).  
5  
6

7  
8  
9 This clustering of chemical variables highlighted five groups corresponding to the sediment  
10 type ("marine", "terrigenous", POC<sub>7</sub> and inorganic trace elements of "natural" or "mixed"  
11 origin) and three groups of anthropogenic contaminants ("inorganic", "organometallic" and  
12 "organic" elements/compounds), resulting in 29 representative chemical variables. The  
13 content ranges, median content and SWA (only for the 23 variables measured on all the  
14 sediments samples) values of these chemical variables are summarized in table 1.  
15  
16  
17  
18  
19  
20  
21

22 Enrichment factors (EF) (Hilton et al., 1985) were calculated for trace metals/metalloids  
23 contaminants in order to distinguish anthropogenic pollution from the natural content of the  
24 sediment as described in Tessier et al. (2011). Briefly, the measured trace elements contents  
25 were normalized to the Al contents of the samples, and then compared with the geochemical  
26 background of the studied area for the same normalized metal content, according to the  
27 following equation:  
28  
29  
30  
31  
32  
33  
34

$$35 \quad EF = \frac{(ME/Al)_{sample}}{(ME/Al)_{background}}$$

36 where Me is the content of the corresponding trace metal/metalloid. The geochemical  
37 background was here estimated as the average of the three lowest calculated Me/Al ratios  
38 from 146 samples originating from surface and deep sediments (including <sup>210</sup>Pb-dated core  
39 sediments corresponding to the lowest contamination records, from Dang et al., 2015a and  
40 Tessier et al., 2011) of Toulon Bay. According to Sutherland (2000), significant enrichment  
41 levels are reached if 5 < EF < 20, and very high enrichment levels if EF > 20.  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51

52 The toxic potential of the sediment was evaluated through comparison of the contents to  
53 effects range-low (ERL) and effects range-medium (ERM) guidelines, initially determined  
54 through toxicity evaluation on benthic invertebrates exposed to single contaminations (Long  
55 et al., 1995). Then, in order to assess the global toxic potential of the multi-contaminated  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

sediments of Toulon Bay, the mean ERM quotient (m-ERM-Q) was calculated as described in Long et al. (1998), according to the following equation:

$$m - ERM - Q = \frac{\sum_{i=1}^n (C_i / ERM_i)}{n}$$

where  $C_i$  is the sediment content of compound  $i$ ,  $ERM_i$  is the ERM for compound  $i$  and  $n$  is the number of compounds.

Local (alpha) diversity indexes (number of terminal restriction fragments (T-RF richness), Pielou's Evenness and Shannon H') were computed using PRIMER 6 software (v6.1.13, Clarke, 1993).

Spatial variations of both chemical composition and microbial communities' T-RF structure (beta-diversity) were analysed through quantitative distance analyses, conducted using PRIMER 6 software. Chemical Gower similarities were computed from median normalized substance contents in order to properly compare minor and major element content<sup>2</sup> variation. Biological Bray-Curtis similarities were computed from  $\text{asin}(\sqrt{x})$  normalized T-RFLP data (Legendre and Legendre, 1998). Differences between samples originating from LiB or LaB were visualized through principal coordinate analysis (PCoA) and tested by one-way permutational multivariate analysis of variance (PERMANOVA) with Rstudio (RStudio Team, 2015, package *vegan*).

Correlations between biological and chemical similarities were assessed by Mantel's test using XLStat (version 2010.3.03, Addinsoft, Paris, France). Other correlations were tested with Spearman's correlation test using Rstudio (RStudio Team, 2015, packages *stats* and *psych*)

Variation partitioning through direct multivariate canonical correspondence analysis (CCA) was used to compare the respective contributions of anthropogenic contaminants and sediment type to the spatial variability of archaeal and bacterial community structure. In order to avoid using multicollinear variables and thus overestimating contributions to the biological variance, covariance analyses were performed and variables presenting variance inflation

factors higher than 10 were excluded from the dataset. Covariance analysis and CCA were performed with Rstudio (RStudio Team, 2015, packages *vegan*, *HH* and *usdm*).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

### 3. RESULTS

#### 3.1. *Abiotic characterization of the sediments*

A set of chemical gradients was observed in surface sediments of Toulon Bay. Groups related to sediment origin and natural and mixed trace elements presented narrow to moderate content gradients (table 1), with Al (figure 2A) and Ca (figure SI-4A) presenting clear terrestrial-marine gradients resulting in an opposite relationship between these 2 elements (figure SI-5). Among inorganic contaminants (ex. Cd, figure 2B), the strongest spatial variations in concentrations was observed for Cu, Hg, Sn and Zn, with about 55- to 900-fold content ranges (table 1). Organo-metals also presented very wide content gradients, with a 13500-fold range for TBT and an approximately 80-fold range for MMHg (table 1). Organic contaminations followed moderate to strong gradients with approximately 15 to 89-fold content ranges for  $\Sigma$ PAH (figure SI-4B), fluoranthene, POC, CB-153 (figure 2C) and  $\Sigma$ PCB, respectively (table 1). Most of the natural or mixed trace elements and inorganic, organometallic or organic contaminants presented similar distribution, their content values being the highest in the northern part of LiB and gradually decreasing toward the south-east and the open sea (for representative examples, see figures 2B, 2C and SI-4A). Tin, POC fluoranthene and  $\Sigma$ PAH did not follow this trend, presenting more random variations (figure SI-4B).

Considering most of these representative physical-chemical parameters together (except for TBT,  $\Sigma$ BT, fluoranthene,  $\Sigma$ PAH, CB-153 and  $\Sigma$ PCB, which were not measured in all of the sampling sites), spatial variations in the global chemical signature of the sediment (*i.e.* considering all 8 of the clusters identified above) were assessed by calculating Gower similarities. This distance analysis demonstrated important variations of sediment chemical composition with reference to the less contaminated sites (two stations located south-east and south-west of Toulon bay, respectively, away from most of the human activities centers), with an offshore-to-coast gradient following the distribution of the main contaminants (figure 2D). Very important shifts of chemical signature were especially recorded in LiB, the most

1 anthropized area, resulting in a very low degree of similarity (decreasing to 28%) compared  
2 with the mouth of the bay. Similar gradients were globally observed when computing Gower  
3 distances for the chemical signatures of the sediment type and the anthropogenic  
4 contaminants, with significant local shifts observed mainly in the northern LiB and near the  
5 coasts of St. Mandrier's peninsula (figure SI-4C and D). Specific distance assessments with  
6 fewer sampling points to include organic contaminants (24 sampling sites) or TBT and  $\Sigma$ BT  
7 (29 sampling sites) confirmed this trend (data not shown).  
8  
9

### 10 **3.2. Contamination levels and potential toxicity**

11 In order to better evaluate sediment contamination by trace metals, Al-normalized  
12 enrichment factors (EF) were calculated to compare the surface sediment concentrations of  
13 Ag, As, Ba, Bi, Cd, Co, Cr, Cu, Hg, Ni, P, Pb, Sb, Sn, Ti and Zn with the geochemical  
14 background of Toulon bay (table 1). Mercury was clearly the most enriched inorganic  
15 contaminant in these sediments, reaching an EF as high as 1200 and presenting very high  
16 enrichment levels (EF > 20) in approximately 80% of the surface sediment area of Toulon  
17 Bay (table 1). A few other inorganic contaminants (Ag, Bi, Cu, Pb, Sb, Sn, and Zn) presented  
18 significant enrichment levels (5 < EF < 20) over 10% to 90% of the surface of Toulon Bay  
19 and very high local enrichment levels, reaching up to 50 and 54 for Sn and Cu, respectively.

20 As detailed in table 1, the highest recorded concentrations of five inorganic contaminants  
21 (As, Cu, Hg, Pb and Zn) and of two organic contaminants (fluoranthene and  $\Sigma$ PCB)  
22 surpassed their ERM values. Moreover, approximately 50% of the surface of Toulon Bay  
23 presented Hg contents overpassing the corresponding ERM. Then, m-ERM-Q was  
24 calculated in order to compare the toxic potential of the different sediment samples. When  
25 considering only trace metals, 26.6% of the whole surface of Toulon bay presented m-ERM-q  
26 values above 0.5, representing a probability of being toxic of approximately 50% or higher.  
27 Moreover, more than 80% of the surface presented m-ERM-Q values above 0.5. According  
28 to spatial variation in metals-based m-ERM-Q, the highest toxic potentials were observed in  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

the northern part of LiB (figure 2E). Including organic pollutants also confirmed this trend (figure SI-4E).

### **3.3. Local microbial diversity variations**

Bacterial T-RF richness ranged between 14 and 40 OTUs. It appeared to be higher in LiB, and decreased gradually through LaB where it reached its minimum before increasing slightly towards the open sea (figure 3A). Archaeal T-RF richness appeared to be smaller with 4 to 13 T-RFs, and varied in a more chaotic way with important changes between very close sampling stations (figure 3B). Shannon indexes calculated from T-RFLP data provided very similar results ( $H'$  ranging from 1.34 to 2.43 for archaea and from 2.60 to 3.52 for bacteria) and were very strongly correlated to richness ( $R^2 = 0.96$  for archaea and 0.94 for bacteria). Evenness demonstrated very low spatial variations (ranging from 0.91 to 0.995 for both bacteria and archaea) and was not correlated with Shannon  $H'$  variations ( $R^2 < 0.1$  for both archaea and bacteria), thus highlighting the importance of richness in bacterial and archaeal Shannon index variations among sediment samples. Hence, local microbial diversity will be represented by genetic richness in further analyses.

Bacterial T-RF richness was positively correlated with parameters related to “terrigenous” elements (Al, Fe), one “natural” trace element (Ni), trace elements of “mixed” origin (As, Cr and S), some anthropogenically enriched toxic trace metals (Ag, Cu, Hg and Zn), organometallic contaminants (MMHg, TBT and  $\Sigma$ BT ), some organic contaminants (CB-153,  $\Sigma$ PCB) and metal-based m-ERM-Q, whereas it was correlated negatively with marine elements (Ca), Co and Ni ( $p < 0.05$ , table 2).

Archaeal genetic richness proved to be linked to fewer environmental parameters: it was also positively correlated with terrigenous elements (Al, Fe), one trace element of mixed origin (Cr), some anthropogenically enriched toxic trace metals (Bi, Cu, Hg, Pb and Zn), organometallic contaminants (MMHg, TBT and  $\Sigma$ BT ) and m-ERM-Q values ( $p < 0.05$ , table 2).



1 Hence, in Toulon Bay, both bacterial and archaeal T-RF richness appeared to be higher in  
2 sediments influenced by terrestrial inputs, enriched in many toxic substances, and thus  
3 presenting a high toxic potential. However, for most of the chemical substances presenting  
4 significant correlations with the T-RF richness of both microbial communities, the correlations  
5 were weaker and less significant for archaea. Particulate organic carbon, Mn and P contents  
6 in the sediments did not appear to be linked with either bacterial or archaeal T-RF richness  
7 ( $p > 0.05$ , table 2)  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17

### 18 **3.4. Spatial variations in microbial community structure**

19 Beta- diversity analysis (PERMANOVA) of the bacterial community highlighted a clear  
20 segregation ( $p = 0.001$ ) of the samples originating, on one hand, from LiB and, on the other  
21 hand, from LaB and towards the open sea (figure 4A). With reference to the less  
22 contaminated sites located outside of Toulon Bay, the bacterial similarity appeared very low  
23 (decreasing to approximately 20%) upon entry into LaB, then increased slightly but varied  
24 strongly again between very close stations westward (figure 3C). Complementary analysis  
25 with reference to the most contaminated site (MIS station, in the northern part of LiB, figure  
26 SI-4F) demonstrated very strong differentiation towards the south of the bay, with low  
27 similarity (decreasing to 50%) between sites distant from barely more than 1km. In contrast,  
28 no evident clustering according to geographical origin ( $p = 0.052$ ) could be observed for the  
29 archaeal community (figure 4B). The archaeal similarities were globally higher and less  
30 variable, with the strongest shifts being observed in LiB and near the mouth eastward (figure  
31 3D).  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

48 Both archaeal and bacterial similarities were negatively correlated with the geographical  
49 distance between sites ( $n = 703$ , archaea:  $r = -0.286$ ,  $p < 10^{-4}$ , bacteria:  $r = -0.285$ ,  $p < 10^{-4}$ ),  
50 and positively correlated to the sediment type's chemical signature ( $n = 703$ , archaea:  $r =$   
51  $0.195$ ,  $p < 10^{-4}$ , bacteria:  $r = 0.209$ ,  $p < 10^{-4}$ ). However, the spatial variation in the global  
52 chemical signature only appeared very weakly correlated with the bacterial similarities ( $n =$   
53  $703$ ,  $r = 0.094$ ,  $p = 0.014$ ) and not significantly correlated with the archaeal similarities ( $n =$   
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

703,  $p > 0.05$ ). The anthropogenic contaminants' chemical signature did not demonstrate any significant correlation with biological similarities ( $n = 703$ ,  $p > 0.05$ ). The same trends were recorded with reduced datasets including BT species ( $n = 276$ ) or organic contaminants ( $n = 171$ ), while archaeal similarities only correlated significantly with the geographic distances between sites (data not shown).

When considering all of the sampling stations ( $n = 38$ ), the combination of sediment type (represented by Ca, Co, As, Cr, Mn, S and POC) and anthropogenic contaminants (Ba, Cd, Hg, P, Sb, Sn and Ti) explained 46% of the bacterial community's structural variation, each of them accounting for 26% on its own. Similarly, this dataset explained 51% of the archaeal community's structural variation, with anthropogenic contaminants explaining a little less than sediment type (26% and 26%, respectively). Including BT species did not significantly increase the contribution of the anthropogenic contaminants, since TBT and  $\Sigma$ BT never explained more than 5% of the variability of the archaeal and bacterial communities' structure, far less than the other contaminants, which contributed up to 9% in the same dataset ( $n = 24$ ). However, the global statistical contribution of the chemical dataset to the biological variation rose to 60% for both bacteria and archaea. Similarly, CB-153 and  $\Sigma$ PCB proved to be the least important statistical contributors to biological variation, with contributions ranging between 4% and 6%, whereas other single contaminants could explain up to 13%. On the contrary, fluoranthene and  $\Sigma$ PAH contributed significantly (between 6% and 10%), and appeared to be among the four best statistical contributors of all the contaminants in the corresponding dataset ( $n = 19$ ). Within this reduced dataset, the global statistical contribution to biological variation reached 85% and 78% for bacteria and archaea, respectively.

#### 4. DISCUSSION

1  
2 Marine coastal areas are subject to various natural and anthropogenic gradients. The benthic  
3 compartment of Toulon Bay illustrates well this general observation with gradients of POC  
4 content, sediment origin and multiple contaminations. In the literature, previous attempts to  
5 understand the contributions of the various factors shaping prokaryotic communities in  
6 different ecosystems mainly focused on the relative contributions of bottom-up and top-down  
7 variables to better understand food-web functioning (e.g. Berdjeb et al., 2011; Salcher, 2013;  
8 Van Wambeke et al., 2009) or on the influence of chemical pollutants through  
9 ecotoxicological approaches (e.g. Besaury et al., 2014; Duran et al., 2015; Kostka et al.,  
10 2011; Paisse et al., 2011; Quillet et al., 2012; Rasmussen and Sorensen, 2001). In this  
11 study, we chose to investigate the influence of the abiotic environment as a whole on  
12 microbial communities by considering together both potentially toxic contaminants and  
13 natural substances from various origins. The usefulness of such a global approach was  
14 demonstrated in this study by the major contribution of the chemical composition of the  
15 sediment in shaping benthic microbial communities of Toulon Bay. Indeed, from  
16 approximately 45% to more than 80% of the variability of bacterial and archaeal community  
17 structure could be statistically explained by a selection of chemical variables. Thus, a  
18 significant part of this variability remained statistically unexplained in our study,  
19 corresponding to some abrupt shifts that were observed in close and chemically similar  
20 sampling sites. This could be partly due to biotic factors, such as predation or viral lysis  
21 pressures, which are well known for their influence on prokaryotic community abundance and  
22 structure (Ben Said et al., 2015; Louati et al., 2013; Ram et al., 2014; Sauret et al., 2015) but  
23 were not investigated in this work.

24  
25 Investigating global chemical signatures also proved to be valuable through the significant  
26 links that were observed between biological and chemical distances. Interestingly, neither the  
27 global chemical distance between the different sampling sites, nor the distance related to  
28 anthropogenic contaminants appeared strongly linked to biological distances, whereas  
29 geographical distances demonstrated stronger links with biological ones. This observation

1 tends to demonstrate that, in this contaminated sediment, local selective pressures are  
2 stronger than the dispersal rate of benthic microbial communities, in spite of the very limited  
3 geographic scale. The positive relationship between biological distances and the chemical  
4 distance related to sediment type (including its marine or terrestrial origin) suggests that  
5 benthic microbial dispersal would be limited to sediment transport processes, which could  
6 promote strong spatial differentiation. However, the high connectivity through the water  
7 column within this marine coastal area, with a mean annual water exchange time of 3.4 days,  
8 makes this hypothesis unlikely (Dufresne et al., 2014). Thereafter, high dispersal but low  
9 survival of benthic microorganisms, because of major chemical shifts between the origin and  
10 end point environments, appears to be a more reliable hypothesis. Future deep taxonomic  
11 investigations will help to test these two hypotheses.

12 Thus, when facing the complexity of chemical variable interactions, such as the antagonistic  
13 effects of nutrients and pollutants, an in-depth geochemical characterization of the  
14 environment appears useful for providing a holistic and detailed chemical signature of the  
15 samples. Then comparing this global chemical signature to other environmental constraints,  
16 such as biotic interactions, appears to be a promising approach to understanding the ecology  
17 of microorganisms and the influence of the human footprint on ecosystem functioning.

18 Another striking result of this study came from multivariate analyses and linear regression  
19 analyses to evaluate the relative contributions of different variable classes to the observed  
20 biological variation. Among the abiotic parameters measured, metallic contamination clearly  
21 appeared to be an influential factor in shaping benthic prokaryote community structure in  
22 Toulon Bay. Indeed, as demonstrated by partial CCA results, at least 26% of the spatial  
23 variation in bacterial and archaeal community structure could be explained by a selection of  
24 trace metal contaminants. The gradient of benthic contamination in Toulon Bay by such  
25 pollutants includes both very low concentrations (*i.e.*, close to the geochemical background  
26 of the region) and dramatic enrichments (*e.g.*, Hg contamination), when compared with other  
27 ecosystems (for a more detailed comparison, see Tessier et al., 2011). Then, considering the

1 panel of potential toxic effects these pollutants are known to have on microorganisms (for  
2 review, see Gadd, 2000; Nies, 1999), observing biological gradients in Toulon Bay appears  
3 logical. These results supported the intention to use bacterial communities as sensitive  
4 indicators of contaminant stress for use as an environmental monitoring tool (Sun et al.,  
5 2012). However, we did not observe any reduction in prokaryotic T-RF richness where trace  
6 metals and organometallic substances were the most enriched. On the contrary, we  
7 observed higher bacterial richness in the north-western bay, the most polluted area. This  
8 observation was confirmed by numerous positive, strong and significant correlations of T-RF  
9 richness with the concentrations of metallic and organometallic pollutants. In the same way,  
10 strong and positive correlations were also observed between T-RF richness and the toxic  
11 potential of the sediment estimated by the m-ERM-Q. However, caution is needed as this  
12 index is based on the sensitivity of macroorganisms rather than that of microorganisms (Long  
13 et al., 1998). Therefore, m-ERM-Q is not well suited for predicting microbial community  
14 responses to chemical contaminations. Whereas metals and organometallic substances are  
15 well known for their toxic effects, their high concentrations were associated with a higher  
16 microbial richness. This striking relation tends to contradict the global erosion of marine  
17 biodiversity caused by anthropogenic inputs in the marine environment (Johnston and  
18 Roberts, 2009). However, the responses of microbial diversity to contaminants are more  
19 complex than those of marine macrobiota and similar observations have been reported in  
20 other marine coastal sediments (Duran et al., 2015; Li et al., 2013; Paissé et al., 2008).

21 One way to explain this contradiction relies on the intermediate disturbance hypothesis  
22 (IDH), which predicts the highest diversity for disturbances of intermediate frequency and  
23 intensity (Connell, 1978). Indeed, although concentrations of the pollutants observed in  
24 Toulon Bay can strongly disturb macroorganisms due to bioaccumulation and lower  
25 reproduction rates, they may induce a lower disturbance to the microbiota that grow faster  
26 and is able to resist and transform metal and organo-metallic substances (Nies, 1999).  
27 Moreover, the most enriched pollutants in the sediments of Toulon Bay are historical  
28 contaminants that have been present in excess for several decades (Pougnnet et al., 2014;

1 Tessier et al., 2011). This stability of the contamination level at the scale of microbial lifetime  
2 could therefore no longer represent a disturbance with respect to the IDH. Indeed, the  
3 bacterial community might have evolved and could now be particularly adapted to such an  
4 environmental stress, as observed in similar ecosystems impacted by metallic pollutions (e.g.  
5 Besaury et al., 2014). In chronically polluted systems, diversity indexes may be a non-  
6 adapted indicator of ecosystem stress as it can recover due to the divergence and  
7 proliferation of tolerant species (Gillan et al., 2005).  
8  
9

10  
11  
12  
13  
14  
15  
16  
17 Organic contaminants appeared to be a less influential environmental stress on benthic  
18 prokaryote communities than metallic and organo-metallic contaminations in Toulon Bay  
19 sediments by statistic disentangling. Indeed, only a few links between organic contamination  
20 and prokaryote T-RF richness were observed in the sediment of Toulon Bay. Moreover, the  
21 contribution of organic contaminants to beta diversity patterns appeared limited. This  
22 observation is quite surprising when considering the relatively high concentrations measured  
23 in comparison to other sites and with reference to sediment quality guidelines (e.g. Gao et  
24 al., 2013; Long et al., 1998; Wolska et al., 2014). Moreover, several studies reported  
25 decreases, or at least very significant shifts, in microbial diversity in response to PAHs or  
26 PCBs contamination (e.g. Correa et al., 2010; Guazzaroni et al., 2013). This could be  
27 explained by the difference of more than one order of magnitude observed between the  
28 respective extents of contamination gradients of metallic and organic pollutants. The  
29 influence of organic contamination could thus be hidden by that of toxic trace metals.  
30 Nevertheless, as for trace metals, organic contaminations appeared to be positively related  
31 to prokaryote T-RF richness in Toulon Bay. Similar observation were previously reported in  
32 other ecosystems (Bengtsson et al., 2013), which could indicate a characteristic feature of  
33 contaminated sites with multiple contaminant types. Indeed, microorganisms are also known  
34 for their ability to grow on organic substrates, and especially PAH. Organic and more or less  
35 biodegradable contamination could thus increase the diversity of carbon resources for  
36 heterotrophic prokaryotes, creating new ecological niches. Thus, in agreement with IDH,  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

organic contamination could partly explain the higher microbial richness found in the north-western part of the bay as previously demonstrated (Bengtsson et al., 2013). Organic contamination could actually counterbalance the degradation of microbial diversity that could occur because of toxic metal enrichment, and finally reduce the extent of disturbance of these microorganisms.

The chemical gradients observed in the sediment of Toulon Bay did not present the same relations with the archaeal and bacterial benthic communities. This observation suggests differential responses of bacteria and archaea to this kind of chemical contamination. The spatial variations of archaeal T-RF richness were weaker than for bacteria, without a significant increase in the most polluted area, and demonstrated fewer and weaker links to the abiotic variables recorded, especially those corresponding to the sediment origin or to biologically relevant substances (POC, S and Mn). Although the low number of archaeal T-RFs detected could raise some methodological concerns, such as partial recovery of the archaeal community, the validation of the method in various previous studies (e.g. Stevenson et al., 2014) tends to invalidate this argument. Moreover, the weak relation between archaeal T-RF richness and the chemical substances measured is counterbalanced by the explanation of 26% to nearly 50% of the spatial variation of archaeal community structure by anthropogenic contaminants, as observed for the bacterial community. This observation could indicate that, whether or not the richness of the community was only poorly affected, its composition was likely more affected. From an ecological point of view, metal resistance in archaea and bacteria involves similar enzymatic responses (Bini, 2010), but archaea are known to predominate in extreme environments, especially where they are exposed to particularly high metal stress as in extremely acidic ecosystems (Baker and Banfield, 2003; Bini, 2010; Edwards et al., 2000). The question arises as to whether archaeal communities could present different and more widespread adaptive strategies than bacterial communities when facing multi-contaminations, especially the combination of organic, organometallic and inorganic contaminants. Further inventories of structural and functional diversity of bacteria

1  
2 and, above all, archaea are now required to explain the different trends we observed in this  
3 study between these communities.  
4  
5

6 Whether inorganic, organometallic or organic, the influence of a pollutant or chemical cocktail  
7 depends on its bioavailability. Although not studied in this work, Dang et al. (2014b, 2015a)  
8 have previously demonstrated that trace metals and metalloids could be highly mobilized  
9 from particles to pore water in surface sediments (0-10 cm), and that this mobilization could  
10 vary throughout the year. This suggests a potentially high bioavailability of inorganic  
11 contaminants in the sediment of Toulon Bay, and tends to support the hypothesis of well-  
12 adapted communities. Thus the observed link between the mobility of inorganic contaminants  
13 and diagenesis activity throughout the seasons (Dang et al., 2014b, 2015a) also suggests  
14 that, in addition to being well adapted, microbial communities could drive the fate of these  
15 contaminants in the sediments of Toulon Bay. Now, investigating microbial community  
16 composition and focusing on functional communities with higher resolution methods under  
17 controlled conditions could help to shed the light on these different interactions between  
18 chemical contamination and microbial communities, as well as on potential cross-interactions  
19 between contaminant types.  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41

## 42 **5. CONCLUSION**

43 Through this study, we have demonstrated the strong influence of the sediments' chemical  
44 signature on marine coastal benthic prokaryotic community diversity. In the sediments of  
45 Toulon Bay, the influence of such a chemical signature appeared to be mainly related to  
46 contamination gradients of trace metal and organo-metallic compounds, which proved to be  
47 strongly related to spatial variation in alpha and beta diversity of bacteria, and to a lesser  
48 extent, of archaea. Although bacterial and archaeal diversity appeared to be linked to this  
49 chemical signature, the major contamination gradients in Toulon Bay surface sediments did  
50 not appear to have resulted in diversity loss. On the contrary, bacterial richness seemed to  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

be higher in the most polluted areas. The underlying ecological adaptive processes and the cross interactions between the multiple pollutant classes now require further clarification.

The current socio-economical context leads more and more scientists to focus on the consequences of human-related inputs for the environment, and especially on the threat they represent to living organisms and natural communities. For instance, it was recently demonstrated that the high Pb levels observed both in farmed and wild mussels from Toulon Bay, mostly originate from Pb sediment contamination, evidencing a clear link between the contaminated abiotic compartment and biotic species (Dang et al., 2015b). However, microorganisms are also well known for their role in the transformation of chemical contaminants, and especially in the geochemistry of toxic trace metals. Our work, in which no apparent toxic effect was observed in spite of very high contamination levels, reminds us how difficult it can be to evaluate which component, *i.e.*, the chemical or the microbiological environment, influences the other to the greatest extent, even in multi-contaminated sites. The microbial contribution to the geochemical cycles of trace metals is still of major interest, and this study illustrates the need for better understanding to fully elucidate the consequences of toxic trace metal inputs to biodiversity. Further in-depth analyses of structural and functional diversities within microbial communities will help in identifying key species and adaptive mechanisms.

## **ACKNOWLEDGEMENTS**

This study was performed in the framework of MerMex-WP3-C3A research program (part of the international IMBER project), CARTOCHIM and CARTOC research programs (funded by "Toulon-Provence-Méditerranée (TPM)", the "Région PACA", and "l'Agence de l'Eau Rhône-Méditerranée et Corse"). The authors wish to thank the French Navy for diving and sampling assistance as well as the LASEM-Cherbourg laboratory (Aline Olivier, Laurent Lacombe) for PAHs and CBs analysis on sediments. We acknowledge the technical assistance of Claire Gassie (EEM) and the Regional Platform for Environmental Microbiology PREMICE supported by the Aquitaine Regional Government Council (France), the urban community of

Pau-Pyrénées (France) and the CNRS. The English grammar and syntax of the manuscript  
have been revised by Proof-Reading-Service.com

## REFERENCES

- Abell, G.C.J., Ross, D.J., Keane, J., Holmes, B.H., Robert, S.S., Keough, M.J., Eyre, B.D., Volkman, J.K., 2014. Niche Differentiation of Ammonia-Oxidising Archaea (AOA) and Bacteria (AOB) in Response to Paper and Pulp Mill Effluent. *Microb. Ecol.* 67, 758–768. doi:10.1007/s00248-014-0376-7
- Baker, B.J., Banfield, J.F., 2003. Microbial communities in acid mine drainage. *FEMS Microbiol. Ecol.* 44, 139–152. doi:10.1016/S0168-6496(03)00028-X
- Bengtsson, G., Törneman, N., De Liphay, J.R., Sørensen, S.J., 2013. Microbial diversity and PAH catabolic genes tracking spatial heterogeneity of PAH concentrations. *Microb. Ecol.* 65, 91–100. doi:10.1007/s00248-012-0112-0
- Ben Said, O., Louati, H., Soltani, A., Preud'homme, H., Cravo-Laureau, C., Got, P., Pringault, O., Aissa, P., Duran, R., 2015. Changes of benthic bacteria and meiofauna assemblages during bio-treatments of anthracene-contaminated sediments from Bizerta lagoon (Tunisia). *Environ. Sci. Pollut. Res. Int.* doi:10.1007/s11356-015-4105-7
- Berdjeb, L., Ghiglione, J.-F., Jacquet, S., 2011. Bottom-up versus top-down control of hypo- and epilimnion free-living bacterial community structures in two neighboring freshwater lakes. *Appl. Environ. Microbiol.* 77, 3591–3599. doi:10.1128/AEM.02739-10
- Besaury, L., Ghiglione, J.-F., Quillet, L., 2014. Abundance, activity, and diversity of archaeal and bacterial communities in both uncontaminated and highly copper-contaminated marine sediments. *Mar. Biotechnol. N. Y.* N 16, 230–242. doi:10.1007/s10126-013-9542-z
- Bini, E., 2010. Archaeal transformation of metals in the environment. *FEMS Microbiol. Ecol.* 73, 1–16. doi:10.1111/j.1574-6941.2010.00876.x
- Burdige, D.J., 2007. Preservation of Organic Matter in Marine Sediments: Controls, Mechanisms, and an Imbalance in Sediment Organic Carbon Budgets? *Chem. Rev.* 107, 467–485. doi:10.1021/cr050347q
- Clarke, K.R., 1993. Non-parametric multivariate analyses of changes in community structure. *Aust. J. Ecol.* 18, 117–143. doi:10.1111/j.1442-9993.1993.tb00438.x
- Colombo, J.C., Cappelletti, N., Lasci, J., Migoya, M.C., Speranza, E., Skorupka, C.N., 2006. Sources, vertical fluxes, and equivalent toxicity of aromatic hydrocarbons in coastal sediments of the Rio de la Plata Estuary, Argentina. *Environ. Sci. Technol.* 40, 734–740. doi:10.1021/es051672y
- Connell, J.H., 1978. Diversity in tropical rain forests and coral reefs. *Science* 199, 1302–1310. doi:10.1126/science.199.4335.1302
- Correa, P.A., Lin, L., Just, C.L., Hu, D., Hornbuckle, K.C., Schnoor, J.L., Van Aken, B., 2010. The effects of individual PCB congeners on the soil bacterial community structure and the abundance of biphenyl dioxygenase genes. *Environ. Int.* 36, 901–906. doi:10.1016/j.envint.2009.07.015
- Cossa, D., Garnier, C., Buscail, R., Elbaz-Poulichet, F., Mikac, N., Patel-Sorrentino, N., Tessier, E., Rigaud, S., Lenoble, V., Gobeil, C., 2014. A Michaelis–Menten type equation for describing methylmercury dependence on inorganic mercury in aquatic sediments. *Biogeochemistry* 119, 35–43. doi:10.1007/s10533-013-9924-3
- Culman, S.W., Bukowski, R., Gauch, H.G., Cadillo-Quiroz, H., Buckley, D.H., 2009. T-REX: software for the processing and analysis of T-RFLP data. *BMC Bioinformatics* 10, 171. doi:10.1186/1471-2105-10-171

- 1 Dang, D.H., Lenoble, V., Durrieu, G., Mullot, J.-U., Mounier, S., Garnier, C., 2014a.  
2 Sedimentary dynamics of coastal organic matter: An assessment of the porewater  
3 size/reactivity model by spectroscopic techniques. *Estuar. Coast. Shelf Sci.* 151, 100–  
4 111. doi:10.1016/j.ecss.2014.10.002
- 5 Dang, D.H., Lenoble, V., Durrieu, G., Omanović, D., Mullot, J.-U., Mounier, S., Garnier, C.,  
6 2015a. Seasonal variations of coastal sedimentary trace metals cycling: insight on the  
7 effect of manganese and iron (oxy)hydroxides, sulphide and organic matter. *Mar.*  
8 *Pollut. Bull.* 92, 113–124. doi:10.1016/j.marpolbul.2014.12.048
- 9 Dang, D.H., Schäfer, J., Brach-Papa, C., Lenoble, V., Durrieu, G., Dutruch, L., Chiffoleau, J.-  
10 F., Gonzalez, J.-L., Blanc, G., Mullot, J.-U., Mounier, S., Garnier, C., 2015b.  
11 Evidencing the Impact of Coastal Contaminated Sediments on Mussels Through Pb  
12 Stable Isotopes Composition. *Environ. Sci. Technol.* 49, 11438–11448.  
13 doi:10.1021/acs.est.5b01893
- 14 Dang, D.H., Tessier, E., Lenoble, V., Durrieu, G., Omanović, D., Mullot, J.-U., Pfeifer, H.-R.,  
15 Mounier, S., Garnier, C., 2014b. Key parameters controlling arsenic dynamics in  
16 coastal sediments: An analytical and modeling approach. *Mar. Chem.* 161, 34–46.  
17 doi:10.1016/j.marchem.2014.02.005
- 18 Daskalakis, K., Oconnor, T., 1995. Distribution of Chemical Concentrations in Us Coastal  
19 and Estuarine Sediment. *Mar. Environ. Res.* 40, 381–398. doi:10.1016/0141-  
20 1136(94)00150-N
- 21 Dufresne, C., Duffa, C., Rey, V., 2014. Wind-forced circulation model and water exchanges  
22 through the channel in the Bay of Toulon. *Ocean Dyn.* 64, 209–224.  
23 doi:10.1007/s10236-013-0676-3
- 24 Duran, R., Bielen, A., Paradžik, T., Gassie, C., Pustijanac, E., Cagnon, C., Hamer, B.,  
25 Vujaklija, D., 2015. Exploring Actinobacteria assemblages in coastal marine  
26 sediments under contrasted Human influences in the West Istria Sea, Croatia.  
27 *Environ. Sci. Pollut. Res. Int.* doi:10.1007/s11356-015-4240-1
- 28 Duran, R., Ranchou-Peyruse, M., Menuet, V., Monperrus, M., Bareille, G., Goni, M.S.,  
29 Salvado, J.C., Amouroux, D., Guyoneaud, R., Donard, O.F.X., Caumette, P., 2008.  
30 Mercury methylation by a microbial community from sediments of the Adour Estuary  
31 (Bay of Biscay, France). *Environ. Pollut.* 156, 951–958.  
32 doi:10.1016/j.envpol.2008.05.007
- 33 Edwards, C.D., Beatty, J.C., Loiselle, J.B.R., Vlassov, K.A., Lefebvre, D.D., 2013. Aerobic  
34 transformation of cadmium through metal sulfide biosynthesis in photosynthetic  
35 microorganisms. *Bmc Microbiol.* 13, 161. doi:10.1186/1471-2180-13-161
- 36 Edwards, K.J., Bond, P.L., Gihring, T.M., Banfield, J.F., 2000. An Archaeal Iron-Oxidizing  
37 Extreme Acidophile Important in Acid Mine Drainage. *Science* 287, 1796–1799.  
38 doi:10.1126/science.287.5459.1796
- 39 Filipkowska, A., Kowalewska, G., Pavoni, B., 2014. Organotin compounds in surface  
40 sediments of the Southern Baltic coastal zone: a study on the main factors for their  
41 accumulation and degradation. *Environ. Sci. Pollut. Res.* 21, 2077–2087.  
42 doi:10.1007/s11356-013-2115-x
- 43 Francis, C.A., Roberts, K.J., Beman, J.M., Santoro, A.E., Oakley, B.B., 2005. Ubiquity and  
44 diversity of ammonia-oxidizing archaea in water columns and sediments of the ocean.  
45 *Proc. Natl. Acad. Sci. U. S. A.* 102, 14683–14688. doi:10.1073/pnas.0506625102
- 46 Frank, K.L., Rogers, D.R., Olins, H.C., Vidoudez, C., Girguis, P.R., 2013. Characterizing the  
47 distribution and rates of microbial sulfate reduction at Middle Valley hydrothermal  
48 vents. *ISME J.* 7, 1391–1401. doi:10.1038/ismej.2013.17
- 49 Gadd, G.M., 2000. Microbial interactions with tributyltin compounds: detoxification,  
50 accumulation, and environmental fate. *Sci. Total Environ., TBT Costs and Benefits- A*  
51 *258*, 119–127. doi:10.1016/S0048-9697(00)00512-X
- 52 Gao, Y., de Brauwere, A., Elskens, M., Croes, K., Baeyens, W., Leermakers, M., 2013.  
53 Evolution of trace metal and organic pollutant concentrations in the Scheldt River  
54 Basin and the Belgian Coastal Zone over the last three decades. *J. Mar. Syst.* 128,  
55 52–61. doi:10.1016/j.jmarsys.2012.04.002
- 56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1 Gauthier, M., Lafay, B., Christen, R., Fernandez, L., Acquaviva, M., Bonin, P., Bertrand, J.,  
2 1992. *Marinobacter-Hydrocarbonoclasticus* Gen-Nov, Sp-Nov, a New, Extremely  
3 Halotolerant, Hydrocarbon-Degrading Marine Bacterium. *Int. J. Syst. Bacteriol.* 42,  
4 568–576.
- 5 Gendron, A., Silverberg, N., Sundby, B., Lebel, J., 1986. Early diagenesis of cadmium and  
6 cobalt in sediments of the Laurentian Trough. *Geochim. Cosmochim. Acta* 50, 741–  
7 747. doi:10.1016/0016-7037(86)90350-9
- 8 Gillan, D.C., Danis, B., Pernet, P., Joly, G., Dubois, P., 2005. Structure of Sediment-  
9 Associated Microbial Communities along a Heavy-Metal Contamination Gradient in  
10 the Marine Environment. *Appl. Environ. Microbiol.* 71, 679–690.  
11 doi:10.1128/AEM.71.2.679-690.2005
- 12 Gillan, D.C., Roosa, S., Kunath, B., Billon, G., Wattiez, R., 2014. The long-term adaptation of  
13 bacterial communities in metal-contaminated sediments: a metaproteogenomic study.  
14 *Environ. Microbiol.* doi:10.1111/1462-2920.12627
- 15 Guazzaroni, M.-E., Herbst, F.-A., Loes, I., Tamames, J., Isabel Pelaez, A., Lopez-Cortes, N.,  
16 Alcaide, M., Del Pozo, M.V., Maria Vieites, J., von Bergen, M., Gallego, J.L.R.,  
17 Bargiela, R., Lopez-Lopez, A., Pieper, D.H., Rossello-Mora, R., Sanchez, J., Seifert,  
18 J., Ferrer, M., 2013. Metaproteogenomic insights beyond bacterial response to  
19 naphthalene exposure and bio-stimulation. *Isme J.* 7, 122–136.  
20 doi:10.1038/ismej.2012.82
- 21 Hilton, J., Davison, W., Ochsenein, U., 1985. A mathematical model for analysis of  
22 sediment core data: Implications for enrichment factor calculations and trace-metal  
23 transport mechanisms. *Chem. Geol.* 48, 281–291. doi:10.1016/0009-2541(85)90053-  
24 1
- 25 Jean, N., Dumont, E., Durrieu, G., Balliau, T., Jamet, J.-L., Personnic, S., Garnier, C., 2012.  
26 Protein expression from zooplankton communities in a metal contaminated. *Mar.*  
27 *Environ. Res.* 80, 12–26. doi:10.1016/j.marenvres.2012.06.004
- 28 Johnston, E.L., Roberts, D.A., 2009. Contaminants reduce the richness and evenness of  
29 marine communities: A review and meta-analysis. *Environ. Pollut.* 157, 1745–1752.  
30 doi:10.1016/j.envpol.2009.02.017
- 31 Jørgensen, B.B., 2000. Bacteria and Marine Biogeochemistry, in: Schulz, P.D.H.D., Zabel,  
32 D.M. (Eds.), *Marine Geochemistry*. Springer Berlin Heidelberg, pp. 173–207.
- 33 Kostka, J.E., Prakash, O., Overholt, W.A., Green, S.J., Freyer, G., Canion, A., Delgardio, J.,  
34 Norton, N., Hazen, T.C., Huettel, M., 2011. Hydrocarbon-Degrading Bacteria and the  
35 Bacterial Community Response in Gulf of Mexico Beach Sands Impacted by the  
36 Deepwater Horizon Oil Spill. *Appl. Environ. Microbiol.* 77, 7962–7974.  
37 doi:10.1128/AEM.05402-11
- 38 Larrose, A., Coynel, A., Schäfer, J., Blanc, G., Massé, L., Maneux, E., 2010. Assessing the  
39 current state of the Gironde Estuary by mapping priority contaminant distribution and  
40 risk potential in surface sediment. *Appl. Geochem.* 25, 1912–1923.  
41 doi:10.1016/j.apgeochem.2010.10.007
- 42 Legendre, P., Legendre, L., 1998. *Numerical Ecology*, Volume 24, (Developments in  
43 Environmental Modelling).
- 44 Levin, L.A., Ekau, W., Gooday, A.J., Jorissen, F., Middelburg, J.J., Naqvi, S.W.A., Neira, C.,  
45 Rabalais, N.N., Zhang, J., 2009. Effects of natural and human-induced hypoxia on  
46 coastal benthos. *Biogeosciences* 6, 2063–2098. doi:10.5194/bg-6-2063-2009
- 47 Li, J., Li, F., Yu, S., Qin, S., Wang, G., 2013. Impacts of mariculture on the diversity of  
48 bacterial communities within intertidal sediments in the Northeast of China. *Microb.*  
49 *Ecol.* 66, 861–870. doi:10.1007/s00248-013-0272-6
- 50 Long, E.R., Field, L.J., MacDonald, D.D., 1998. Predicting toxicity in marine sediments with  
51 numerical sediment quality guidelines. *Environ. Toxicol. Chem.* 17, 714–727.  
52 doi:10.1002/etc.5620170428
- 53 Long, E.R., Macdonald, D.D., Smith, S.L., Calder, F.D., 1995. Incidence of adverse biological  
54 effects within ranges of chemical concentrations in marine and estuarine sediments.  
55 *Environ. Manage.* 19, 81–97. doi:10.1007/BF02472006

- 1 Louati, H., Ben Said, O., Soltani, A., Got, P., Mahmoudi, E., Cravo-Laureau, C., duran, R.,  
2 Aissa, P., Pringault, O., 2013. The roles of biological interactions and pollutant  
3 contamination in shaping microbial benthic community structure. *Chemosphere* 93,  
4 2535–2546. doi:10.1016/j.chemosphere.2013.09.069
- 5 Nicolau, R., Lucas, Y., Merdy, P., Raynaud, M., 2012. Base flow and stormwater net fluxes of  
6 carbon and trace metals to the Mediterranean sea by an urbanized small river. *Water*  
7 *Res.* 46, 6625–6637. doi:10.1016/j.watres.2012.01.031
- 8 Nies, D.H., 1999. Microbial heavy-metal resistance. *Appl. Microbiol. Biotechnol.* 51, 730–750.
- 9 Ortega-Retuerta, E., Joux, F., Jeffrey, W.H., Ghiglione, J.F., 2013. Spatial variability of  
10 particle-attached and free-living bacterial diversity in surface waters from the  
11 Mackenzie River to the Beaufort Sea (Canadian Arctic). *Biogeosciences* 10, 2747–  
12 2759. doi:10.5194/bg-10-2747-2013
- 13 Oursel, B., Garnier, C., Durrieu, G., Mounier, S., Omanović, D., Lucas, Y., 2013. Dynamics  
14 and fates of trace metals chronically input in a Mediterranean coastal zone impacted  
15 by a large urban area. *Mar. Pollut. Bull.* 69, 137–149.  
16 doi:10.1016/j.marpolbul.2013.01.023
- 17 Oursel, B., Garnier, C., Zebracki, M., Durrieu, G., Pairaud, I., Omanović, D., Cossa, D.,  
18 Lucas, Y., 2014. Flood inputs in a Mediterranean coastal zone impacted by a large  
19 urban area: Dynamic and fate of trace metals. *Mar. Chem., Estuarine*  
20 *Biogeochemistry* 167, 44–56. doi:10.1016/j.marchem.2014.08.005
- 21 Paissé, S., Coulon, F., Goñi-Urriza, M., Peperzak, L., McGenity, T.J., Duran, R., 2008.  
22 Structure of bacterial communities along a hydrocarbon contamination gradient in a  
23 coastal sediment. *FEMS Microbiol. Ecol.* 66, 295–305. doi:10.1111/j.1574-  
24 6941.2008.00589.x
- 25 Paise, S., Duran, R., Coulon, F., Goñi-Urriza, M., 2011. Are alkane hydroxylase genes  
26 (alkB) relevant to assess petroleum bioremediation processes in chronically polluted  
27 coastal sediments? *Appl. Microbiol. Biotechnol.* 92, 835–844. doi:10.1007/s00253-  
28 011-3381-5
- 29 Parks, J.M., Johs, A., Podar, M., Bridou, R., Hurt, R.A., Smith, S.D., Tomanicek, S.J., Qian,  
30 Y., Brown, S.D., Brandt, C.C., Palumbo, A.V., Smith, J.C., Wall, J.D., Elias, D.A.,  
31 Liang, L., 2013. The Genetic Basis for Bacterial Mercury Methylation. *Science* 339,  
32 1332–1335. doi:10.1126/science.1230667
- 33 Pougnet, F., Schäfer, J., Dutruch, L., Garnier, C., Tessier, E., Dang, D.H., Lancelour, L.,  
34 Mullot, J.-U., Lenoble, V., Blanc, G., 2014. Sources and historical record of tin and  
35 butyl-tin species in a Mediterranean bay (Toulon Bay, France). *Environ. Sci. Pollut.*  
36 *Res. Int.* 21, 6640–6651. doi:10.1007/s11356-014-2576-6
- 37 Quillet, L., Besaury, L., Popova, M., Paissé, S., Deloffre, J., Ouddane, B., 2012. Abundance,  
38 diversity and activity of sulfate-reducing prokaryotes in heavy metal-contaminated  
39 sediment from a salt marsh in the Medway Estuary (UK). *Mar. Biotechnol. N. Y. N* 14,  
40 363–381. doi:10.1007/s10126-011-9420-5
- 41 Ram, A.S.P., Palesse, S., Colombet, J., Thouvenot, A., Sime-Ngando, T., 2014. The relative  
42 importance of viral lysis and nanoflagellate grazing for prokaryote mortality in  
43 temperate lakes. *Freshw. Biol.* 59, 300–311. doi:10.1111/fwb.12265
- 44 Raskin, L., Stromley, J.M., Rittmann, B.E., Stahl, D.A., 1994. Group-specific 16S rRNA  
45 hybridization probes to describe natural communities of methanogens. *Appl. Environ.*  
46 *Microbiol.* 60, 1232–1240.
- 47 Rasmussen, L.D., Sorensen, S.J., 2001. Effects of mercury contamination on the culturable  
48 heterotrophic, functional and genetic diversity of the bacterial community in soil. *Fems*  
49 *Microbiol. Ecol.* 36, 1–9. doi:10.1016/S0168-6496(01)00111-8
- 50 Rensing, C., Fan, B., Sharma, R., Mitra, B., Rosen, B.P., 2000. CopA: An Escherichia coli  
51 Cu(I)-translocating P-type ATPase. *Proc. Natl. Acad. Sci. U. S. A.* 97, 652–656.
- 52 RStudio Team, 2015. RStudio: Integrated Development for R. RStudio Inc Boston MA URL  
53 [Httpwww RStudio Comide](http://www.RStudio.com).
- 54 Salcher, M.M., 2013. Same same but different: ecological niche partitioning of planktonic  
55 freshwater prokaryotes. *J. Limnol.* 73.

- 1 Sauret, C., Tedetti, M., Guigue, C., Dumas, C., Lami, R., Pujo-Pay, M., Conan, P., Goutx, M.,  
2 Ghiglione, J.-F., 2015. Influence of PAHs among other coastal environmental  
3 variables on total and PAH-degrading bacterial communities. *Environ. Sci. Pollut.*  
4 *Res. Int.* doi:10.1007/s11356-015-4768-0
- 5 Scott, N.M., Hess, M., Bouskill, N.J., Mason, O.U., Jansson, J.K., Gilbert, J.A., 2014. The  
6 microbial nitrogen cycling potential is impacted by polyaromatic hydrocarbon pollution  
7 of marine sediments. *Aquat. Microbiol.* 5, 108. doi:10.3389/fmicb.2014.00108
- 8 Sklar, F.H., Browder, J.A., 1998. Coastal environmental impacts brought about by alterations  
9 to freshwater flow in the Gulf of Mexico. *Environ. Manage.* 22, 547–562.  
10 doi:10.1007/s002679900127
- 11 Stahl D, Amann R, n.d. Development and application of nucleic acid probes, in: *Nucleic Acid*  
12 *Techniques in Bacterial Systematics.* John Wiley & Sons, Chichester, England, pp.  
13 205–248.
- 14 Stevenson, B.A., Hunter, D.W.F., Rhodes, P.L., 2014. Temporal and seasonal change in  
15 microbial community structure of an undisturbed, disturbed, and carbon-amended  
16 pasture soil. *Soil Biol. Biochem.* 75, 175–185. doi:10.1016/j.soilbio.2014.04.010
- 17 Sun, M.Y., Dafforn, K.A., Brown, M.V., Johnston, E.L., 2012. Bacterial communities are  
18 sensitive indicators of contaminant stress. *Mar. Pollut. Bull.* 64, 1029–1038.  
19 doi:10.1016/j.marpolbul.2012.01.035
- 20 Sutherland, R.A., 2000. Bed sediment-associated trace metals in an urban stream, Oahu,  
21 Hawaii. *Environ. Geol.* 39, 611–627. doi:10.1007/s002540050473
- 22 Tessier, E., Garnier, C., Mullot, J.-U., Lenoble, V., Arnaud, M., Raynaud, M., Mounier, S.,  
23 2011. Study of the spatial and historical distribution of sediment inorganic  
24 contamination in the Toulon bay (France). *Mar. Pollut. Bull.* 62, 2075–2086.  
25 doi:10.1016/j.marpolbul.2011.07.022
- 26 Troussellier, M., Schafer, H., Batailler, N., Bernard, L., Courties, C., Lebaron, P., Muyzer, G.,  
27 Servais, P., Vives-Rego, J., 2002. Bacterial activity and genetic richness along an  
28 estuarine gradient (Rhône River plume, France). *Aquat. Microb. Ecol.* 28, 13–24.  
29 doi:10.3354/ame028013
- 30 Van Wambeke, F., Ghiglione, J.-F., Nedoma, J., Mével, G., Raimbault, P., 2009. Bottom up  
31 effects on bacterioplankton growth and composition during summer-autumn transition  
32 in the open NW Mediterranean Sea. *Biogeosciences* 6, 705–720. doi:10.5194/bg-6-  
33 705-2009
- 34 Waterkeyn, A., Grillas, P., Vanschoenwinkel, B., Brendonck, L., 2008. Invertebrate  
35 community patterns in Mediterranean temporary wetlands along hydroperiod and  
36 salinity gradients. *Freshw. Biol.* 53, 1808–1822. doi:10.1111/j.1365-  
37 2427.2008.02005.x
- 38 Winemiller, K., Leslie, M., 1992. Fish Assemblages Across a Complex, Tropical Fresh-Water  
39 Marine Ecotone. *Environ. Biol. Fishes* 34, 29–50. doi:10.1007/BF00004783
- 40 Wolska, L., Mechlińska, A., Rogowska, J., Namieśnik, J., 2014. Polychlorinated biphenyls  
41 (PCBs) in bottom sediments: Identification of sources. *Chemosphere* 111, 151–156.  
42 doi:10.1016/j.chemosphere.2014.03.025
- 43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Table1

[Click here to download Table: table 1.docx](#)**Table 1.** Chemical characterization of the Toulon bay surface sediments.

Variable groups (unit)	Chemical parameters	N	Content range (median)	SWA content	EF range (median)	Surface (%) per EF level				ERL ( $\mu\text{g g}^{-1}$ )	ERM	Surface (%) per ER level			
						EF < 1.5	1.5 < EF < 5	5 < EF < 20	EF > 20			Content < ERL	ERL < content < ERM	Content > ERM	
Marine ( $\mu\text{g.g}^{-1}$ )	Ca	54	84000 - 320000 (140000)	191000	-	-	-	-	-	-	-	-	-	-	-
Terrigenous ( $\mu\text{g.g}^{-1}$ )	Al	54	4700 - 57000 (29000)	22000	-	-	-	-	-	-	-	-	-	-	-
	Fe	54	5400 - 35000 (19000)	16000	-	-	-	-	-	-	-	-	-	-	-
Natural ( $\mu\text{g.g}^{-1}$ )	Co	54	1.9 - 10 (6.2)	5.56	1 - 6.5 (1.6)	15.8	83.7	0.5	0.0	-	-	-	-	-	-
	Ni	54	5.1 - 40 (22)	17.7	1 - 1.9 (1.3)	91.4	8.6	0.0	0.0	20.9	51.6	65.0	35.0	0.0	-
Mixed ( $\mu\text{g.g}^{-1}$ )	As	54	6.2 - 80 (21)	19.8	0.7 - 21 (2.1)	2.2	92.1	5.7	0.0	8.2	70	1.3	98.5	0.1	-
	Cr	54	12 - 95 (38)	31.9	1.5 - 3.9 (2)	0.9	99.1	0.0	0.0	81	370	99.5	0.5	0.0	-
	Mn	54	99 - 930 (150)	185	-	-	-	-	-	-	-	-	-	-	-
	S	54	2900 - 17000 (6100)	5470	-	-	-	-	-	-	-	-	-	-	-
Inorganic contaminants ( $\mu\text{g.g}^{-1}$ )	Ag	54	0.061 - 2.8 (0.32)	0.31	0.9 - 22 (3.4)	1.6	87.4	11.0	0.1	1.0	3.7	94.4	5.7	0.0	-
	Ba	54	30 - 480 (150)	134	1.7 - 5.2 (2.6)	0.0	99.7	0.3	0.0	-	-	-	-	-	-
	Bi	54	0.17 - 4.6 (0.82)	0.63	1.5 - 25 (5.5)	0.0	48.4	51.5	0.1	-	-	-	-	-	-
	Cd	54	0.07 - 1.7 (0.21)	0.202	1.2 - 14 (2.3)	2.3	88.2	9.6	0.0	1.2	9.6	99.6	0.4	0.0	-
	Cu	54	5.6 - 730 (65)	60.5	1 - 54 (6.4)	1.3	50.3	46.6	1.7	34	270	43.3	54.8	1.9	-
	Hg	54	0.03 - 27 (1.6)	1.54	1.7 - 1200 (77)	0.0	0.7	20.3	79.1	0.15	0.71	11.6	40.7	47.8	-
	P	54	290 - 930 (540)	474	2.1 - 16 (2.9)	0.0	80.8	19.2	0.0	-	-	-	-	-	-
	Pb	54	15 - 470 (99)	77	1.6 - 38 (9.6)	0.0	9.8	89.6	0.6	46.7	218	33.9	64.6	1.5	-
	Sb	54	0.18 - 5.5 (0.62)	0.786	1.8 - 44 (4)	0.0	47.9	50.1	2.1	-	-	-	-	-	-
	Sn	54	0.94 - 120 (10)	12.9	1.3 - 50 (5.2)	2.6	37.7	55.8	4.0	-	-	-	-	-	-
	Ti	54	91 - 640 (370)	320	1.1 - 3.4 (1.7)	21.4	78.6	0.0	0.0	-	-	-	-	-	-
	Zn	54	27 - 1500 (150)	136	1.7 - 31 (4.5)	0.0	77.1	22.6	0.4	150	410	68.6	28.4	3.1	-
Organometallic contaminants ( $\text{ng.g}^{-1}$ )	MMHg	54	0.18 - 15 (2.5)	1.99	-	-	-	-	-	-	-	-	-	-	-
	TBT	29	0.2 - 2700 (24)	-	-	-	-	-	-	-	-	-	-	-	-
	$\Sigma$ BT	29	0.86 - 4500 (140)	-	-	-	-	-	-	-	-	-	-	-	-
Organic contaminants ( $\mu\text{g.g}^{-1}$ )	CB-153	25	0.013 - 0.42 (0.05)	-	-	-	-	-	-	-	-	-	-	-	-
	$\Sigma$ CB	25	0.018 - 1.6 (0.11)	-	-	-	-	-	-	0.023	0.18	-	-	-	-
	fluoranthene	25	0.34 - 6.2 (2.1)	-	-	-	-	-	-	0.60	5.1	-	-	-	-





**Table 2.** Spearman's correlation tests between archaeal or bacterial T-RF richness and chemical variables. Numbers represent  $\rho$  values of statistically significant correlations. n=38 for most of the correlation analyses, except for BTs and organic compounds for which n = 24 and 19, respectively. ns: non significant. \*:  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\* $p < 0.001$ .

Variable groups	Chemical parameters	Archaea	Bacteria
Marine	<i>Ca</i>	ns	-0.38*
Terrigenous	<i>Al</i>	0.35*	0.54***
	<i>Fe</i>	0.36*	0.48**
Natural	<i>Co</i>	ns	ns
	<i>Ni</i>	ns	0.35*
Mixed	<i>As</i>	ns	0.38*
	<i>Cr</i>	0.33*	0.54***
	<i>Mn</i>	ns	ns
	<i>S</i>	ns	0.57***
Inorganic contaminants (EF)	<i>Ag</i>	0.41*	0.70***
	<i>Ba</i>	ns	ns
	<i>Bi</i>	0.50**	ns
	<i>Cd</i>	ns	ns
	<i>Cu</i>	0.39*	0.55***
	<i>Hg</i>	0.41*	0.51**
	<i>P</i>	ns	ns
	<i>Pb</i>	0.45**	ns
	<i>Sb</i>	ns	ns

	<i>Sn</i>	ns	ns
	<i>Ti</i>	ns	ns
	<i>Zn</i>	0.44**	0.48**
Organometallic contaminants	<i>MMHg</i>	0.38*	0.59***
	<i>TBT</i>	0.06*	0.73***
	$\Sigma BT$	0.47*	0.73***
Organic contaminants	<i>Fluoranthene</i>	ns	ns
	$\Sigma PAH$	ns	ns
	<i>CB-153</i>	ns	0.47*
	$\Sigma PCB$	ns	0.47*
Organic matter	<i>POC</i>	ns	ns
Sediment potential toxicity	<i>Metal based m-ERM-Q</i>	0.42**	0.55***
	<i>m-ERM-Q with PAHs</i>	0.50*	ns

## FIGURE CAPTIONS


**Figure 1.** Sampling stations and main anthropogenic activities in Toulon bay. *Dots represents the specific sites where organic contaminants were quantified, small grey squares represent the sampling sites with butyl-tin species measurements and large open squares represent the sites from which samples were analyzed through T-RFLP.*


**Figure 2.** Examples of benthic abiotic spatial gradients in the Toulon bay. **A, B, C:** contents of Al, Cd and CB-153, respectively. **D:** Gower similarity, the white arrows represent the reference sites. **E:** *m-ERM-Q* calculated with toxic metals' concentrations. Scales were drawn according to deciles of the respective datasets.


**Figure 3.** Spatial variations of prokaryotic richness and community structure in surface sediments of Toulon bay, calculated from t-RFLP data. **A and B:** bacterial and archaeal richnesses, respectively. **C and D:** Bray-Curtis similarities of T-RF structure within bacterial and archaeal communities, respectively. Arrows design the reference sites used for representing similarities.


**Figure 4.** PCoA of bacterial (**A**) and archaeal (**B**) communities using Bray-Curtis distances, calculated from normalized tRFLP data. *LiB: Little Bay, LaB: Large Bay*


Figure1  
[Click here to download high resolution image](#)




















**B**







**D**









**Supplementary material for on-line publication only**

**[Click here to download Supplementary material for on-line publication only: Misson et al.\\_SI.docx](#)**